

# REPORT DOCUMENTATION PAGE

Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE Technical Papers	3. DATES COVERED (From - To)		
4. TITLE AND SUBTITLE			5a. CONTRACT NUMBER	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)			5d. PROJECT NUMBER 2303	
			5e. TASK NUMBER M2C8	
			5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT	
Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT				
Approved for public release; distribution unlimited.				
13. SUPPLEMENTARY NOTES				
14. ABSTRACT				
1121 039				
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Leilani Richardson
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	A	19b. TELEPHONE NUMBER (include area code) (661) 275-5015

62

Separate items are enclosed

2-1998-097

MEMORANDUM FOR IN-HOUSE PUBLICATIONS

FROM: PROI (TI) (STINFO)

30 Apr 98

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-1998-097  
**Mario Fajardo and Simon Tam "Solid Parahydrogen" HEDM Conference Presentation (Statement A)**

HIGH ENERGY DENSITY MATTER CONTRACTORS CONFERENCE  
Monterey, CA 20-22 May 1998.

# Solid Parahydrogen

Mario E. Fajardo and Simon Tam  
US Air Force Research Laboratory, Propulsion Directorate  
AFRL/PRSP, Bldg. 8451, Edwards AFB, CA 93524-7680  
[Mario\\_Fajardo@ple.af.mil](mailto:Mario_Fajardo@ple.af.mil)

## Introduction

Atom Doped Cryogenic Solid Propellants  
Laser Ablation/Matrix Isolation Spectroscopy  
Rapid Vapor Deposition of Thick Transparent pH<sub>2</sub> Solids!  
ortho/para Hydrogen Converter

## Research Update

Microscopic Structure of Rapidly Deposited pH<sub>2</sub> Solids  
IR, Raman, x-polarizers  
MIS Spectroscopy in Doped pH<sub>2</sub> and oD<sub>2</sub> Samples  
“High Resolution” IR Absorption Measurements  
CH<sub>4</sub>/pH<sub>2</sub>, CH<sub>3</sub>OH/pH<sub>2</sub>  
Application: CO/pH<sub>2</sub> -- a Molecular Thermometer  
Photochemistry (guest-host reactions)  
O<sub>2</sub>/pH<sub>2</sub>, B<sub>2</sub>H<sub>6</sub>/oD<sub>2</sub>  
Photodynamics (LIF and photobleaching)  
B/pH<sub>2</sub>, B/Ne, Na/pH<sub>2</sub>, Na/Ne

## Conclusions and Future Directions

High Resolution IR Spectroscopy in pH<sub>2</sub> Hosts

DISTRIBUTION STATEMENT A  
Approved for Public Release  
Distribution Unlimited

20021121 039

# High Energy Density Matter (HEDM)

## Cryosolid Propellants

### Objectives

- \* Trap 5% molar concentration of energetic additives in solid hydrogen.
- \* Demonstrate size-scaleable sample production method.

### Payoffs

#### Increased Specific Impulse

$$I_{sp} \propto \sqrt{\Delta H_{sp}}$$

$$\begin{aligned} \text{LOX/LH}_2 : I_{sp} &= 390 \text{ s} \\ 5\% \text{ B/H}_2 + \text{LOX} : I_{sp} &= 500 \text{ s (+30%)*} \end{aligned}$$

\* calculated for  $P_{\text{chamber}} = 1000 \text{ PSIA}$ ,  $P_{\text{exhaust}} = 14.7 \text{ PSIA}$

#### Greater Propellant Density

$$\text{liquid H}_2 @ 20 \text{ K} : \rho = 0.070 \text{ g/cm}^3$$


$$\text{solid H}_2 @ 2 \text{ K} : \rho = 0.087 \text{ g/cm}^3 (+25\%)$$

$$50/50 \text{ liquid He/solid H}_2 : \rho = 0.105 \text{ g/cm}^3 (+50\%)$$


# Scientific/Technological Motivations

<u>Issue</u>	<u>Scientific Motivation</u>	<u>Tech. Application</u>
Chemical stability of M/pH <sub>2</sub> samples	Chemical reactivity @ low T (1-10 K) existence of small reaction barriers in M + H <sub>2</sub> reaction matrix host effects	Identify candidate M's
Microscopic model of sample deposition process	Molecular dynamics of "simple" condensed phase systems (models for more complicated chemistry)	Maximize [M]
Simulation of M/RGS and M/pH <sub>2</sub> spectroscopy	Spectroscopy in condensed phases spectrum $\leftrightarrow$ structure/fluctuations	Measure [M] and determine fuel p
Diffusion/recombination of M's	Diffusion in "classical" and "quantum" solids	Determine thermal stability of M/H <sub>2</sub> fuel
Maximum attainable [M]	Limits of chemical energy storage	fuel performance

# Experimental Diagram



$\text{Li}/\text{H}_2$   $T=3\text{K}$



M.E. RAJARAO, J. Chem. Phys. 98, 110 (1993).

# Optical Scattering in Solid Hydrogen

## Crystal Growing and Quality (p. 81)


“There is a considerable art to growing hydrogen crystals of high quality. Good crystals are always grown slowly from the melt; a rapid freeze from the gas produces snow.”

## Crystallite Light Scattering (p. 83)

“The reason that a good hydrogen crystal is so hard to see is its low refractive index...an estimated 1.16!


Yet a 1 mm-thick layer of hydrogen crystallites can be a completely opaque brown-black.”

P.C. Souers,  
Hydrogen Properties for Fusion Energy  
(UC Press, Berkeley, 1986).




S. Tam + M. E. Fajardo, unpublished.


B / H<sub>2</sub> raw data (c 1993)



# Ortho and Para Hydrogen




(9) — 4



I.F. Silvera,  
Rev. Mod. Phys. **52**, 393 (1980).


# Ortho/Para Hydrogen Converter Bobbin



1/8 inch OD by 1.5 m long copper tube packed with 1.4 g of APACHI catalyst, coiled around bobbin and potted in place with metal-filled thermally conductive epoxy.

Catalyst is activated by heating to 150 °C under vacuum, then with a slow flow of H<sub>2</sub> gas.

# UV Transmission of 1 mm Thick B/pH<sub>2</sub> Sample




# IR Absorption of 6 mm Thick Parahydrogen Solid

$T = 2 \text{ K.}$

Non-observation of the  $Q_1(0)$  transition ( $4153 \text{ cm}^{-1}$ ) demonstrates the absence of  $\text{OH}_2$  impurities, and that the microscopic structure is not amorphous or porous.

Observation of  $S_1(0)$  transition demonstrates the absence of inversion symmetry for some  $\text{H}_2$  molecular environments.

[J. van Kranendonk and H.P. Gush, Phys. Lett. **1**, 22 (1962)]



# Raman Spectra of 4.5 and 6 mm Thick Parahydrogen Solids

Mixed hcp/fcc as-deposited structure, anneals to hcp;  
compare with:


G.W. Collins, et al.,  
Phys. Rev. B **53**, 102 (1996).

(d) sample in (c) warmed to  
4.5 K.

(c) 4.5 mm sample as deposited  
at 3.3 K ( $\Phi = 290$  mmol/hr).

(b) sample in (a) warmed to  
4.5 K.

(a) 6 mm sample as deposited at  
3.1 K ( $\Phi = 200$  mmol/hr).



# Infrared spectroscopic study of rovibrational states of methane trapped in parahydrogen crystal

Takamasa Momose,<sup>a)</sup> Masaaki Miki, Tomonari Wakabayashi,  
and Tadamasa Shida

*Department of Chemistry, Graduate School of Science, Kyoto University, Kyoto 606-01, Japan*

Man-Chor Chan,<sup>b)</sup> Steven S. Lee, and Takeshi Oka

*Department of Chemistry, The University of Chicago, Chicago, Illinois 60637*

(Received 23 June 1997; accepted 15 August 1997)


The  $\nu_3$  and  $\nu_4$  vibrational transitions of methane trapped in solid parahydrogen have been observed by using Fourier transform infrared and high resolution laser spectroscopy. The observed spectrum is interpreted in terms of rovibrational states of the spherical rotor which are subjected to the crystal field splitting. The  $\nu_4$  band shows extremely sharp lines of a width of  $\sim 0.003 \text{ cm}^{-1}$ , while the  $\nu_3$  band exhibits broader lines of a width of  $1 \text{ cm}^{-1}$ . The infrared selection rules derived from an extended group theory to take into account the field effect are consistent with the observed spectra. The intermolecular interaction and the field effect in solid parahydrogen are analyzed quantitatively.  
© 1997 American Institute of Physics. [S0021-9606(97)04743-0]

JCP 107, 7707 (1997)


110 PPM  $\text{CH}_4/\text{pH}_2$ ,  $d = 1.2 \text{ mm}$

st2320a: as dep.  $T=2 \text{ K}$


b: 4.4K, c: 3.0K, d: 2.2K




ST2117A MeOH:pH2 1:560 as deposited at 2 K




## Deposition Cartoon




# Experimental Diagram – Sample Deposition



# IR Absorptions of CO/pH<sub>2</sub>



# Intensity of $2135 \text{ cm}^{-1}$ band vs. Temperature



0.9 mm CO/pH<sub>2</sub> layer  
deposited on top of 1.9 mm  
pure pH<sub>2</sub> layer.


Integrated intensity of  
 $2135 \text{ cm}^{-1}$  band normalized to  
integral over entire CO band  
for as-deposited sample.

Irreversible shift upon  
annealing attributed to  
 $\text{fcc} \rightarrow \text{hcp}$  conversion  
(introduces T measurement  
error  $\approx +0.2 \text{ K}$ ).


**ST2207**

temperature (K)


## “Van’t Hoff Plot”




st2345, diborane/oD<sub>2</sub> (20  $\mu\text{m}$ )




B/pH<sub>2</sub> LIF  
 $\lambda_{\text{exc}} = 217 \text{ nm, } 125 \mu\text{J/pulse}$




B/pH<sub>2</sub> LIF  
 $\lambda_{\text{exc}} = 207, 210, 217, \text{ and } 220 \text{ nm}$




## B/pH<sub>2</sub> UV absorption and photobleaching




## B/pH<sub>2</sub> UV photobleached lineshape




## B/pH<sub>2</sub> LIF Cartoon




$\nu_{17}$   $\text{B}_2\text{H}_6$  in  $\text{B}/\text{pH}_2$  solid  
as deposited and photobleached




## B/Ne UV absorption




B/Ne LIF  
 $\lambda_{\text{exc}} = 210, 213, 215, 217, 219, 220, 221 \text{ nm}$




Na/Ne LIF  
 $\lambda_{\text{exc}} = 535, 540, 545, 550, 555, 560, 565, 570 \text{ nm}$



Na/pH<sub>2</sub> transmission spectrum  
as deposited and photobleached



Na/pH<sub>2</sub> photobleaching  
60 to 90  $\mu$ J/pulse,  $d_{\text{spot}} \approx 4$  mm




## Conclusions

- \* As-deposited pH<sub>2</sub> samples are mixed hcp/fcc close-packed solids that transform to hcp upon annealing to  $\approx 5$  K.
- \* Demonstrated trapping of various dopant atoms, molecules, and ionic species using conventional matrix isolation sources.
- \* Some dopant IR absorption bands show unresolved structure even at 0.1 cm<sup>-1</sup> resolution.
- \* Measured  $\sim 1$  K temperature rises in  $\sim 0.1$  cm thick samples subjected to 10 mW/cm<sup>2</sup> heat loads during deposition; estimated thermal conductivity of rapid vapor deposited pH<sub>2</sub> is  $\sim 1$  mW/cm-K.
- \* Observed guest-host photochemistry during attempts to produce energetic dopants via 193 nm irradiation.
- \* Observed LIF of B atoms in solid pH<sub>2</sub>. Emission lineshape is independent of excitation wavelength. Photobleaching results in uniform changes to absorption lineshape.
- \* LIF of Na atoms in solid pH<sub>2</sub> is at least four orders of magnitude weaker than LIF of Na atoms in solid Ne. Photobleaching effect depends on excitation wavelength and produces varying changes to absorption lineshape.

## Future Directions


- \* Bruker IFS120HR on order.
- \* High resolution IR absorption ( $0.003\text{ cm}^{-1}$ ) and Raman ( $0.05\text{ cm}^{-1}$ ) spectroscopies:
  - unresolved rotational (hindered rotor?) structure in presently available spectra.
  - determine inhomogeneities in dopant environment:
    - hcp/fcc vs. random stacked for as-deposited dopant-dopant interactions (clusters).
- \* Analysis and simulation of IR spectra:
  - dopant absorptions
    - CO/pH<sub>2</sub> "crystal field" model
      - (collaboration with T. Momose, Kyoto U)
      - direct simulation given dopant-H<sub>2</sub> potentials.
    - induced H<sub>2</sub> absorptions
- \* Measure fluorescence decay of "B atom" LiF.
- \* Work to increase dopant concentrations from 0.1 to 1% levels; demonstrate useful energy storage.

# Substrate and Bulk Hydrogen Temperatures During Deposition



# 1-D Heat Transfer

$$T_{lo} \quad T_{hi}$$



$$\dot{Q}/A = -\kappa \Delta T/\Delta x$$

$$\Delta T = T_{hi} - T_{lo}$$

$\kappa$  is the thermal conductivity

units:

$$\dot{Q}/A \text{ (mW/cm}^2\text{)}$$

$$\Delta T \text{ (K)}$$


$$\Delta x \text{ (cm)}$$

$$\kappa \text{ (mW/cm-K)}$$

$$\Delta x$$

note: 1 mW/cm-K = 0.1 W/m-K

# Calculated Heat Flux vs. pH<sub>2</sub> Deposition Rate



# Thermal Conductivity of Rapid Vapor Deposited pH<sub>2</sub>

Expt.	$[T_{co} - T_B]$ (K)	$\Delta x$ (cm)	$\dot{Q}/A$ (mW/cm <sup>2</sup> )	$\kappa$ (mW/cm-K)	$\kappa$ (W/m-K)
ST2212	0.61	0.12	9.2	1.8	0.18
ST2207	1.19	0.22	13.8	2.6	0.26
ST2217	1.74	0.25	16.6	2.4	0.24
Expt.	$[T_{co} - T_B]$ (K)	$\Delta x$ (cm)	$\dot{Q}/A$ (mW/cm <sup>2</sup> )	$\kappa$ (mW/cm-K)	$\kappa$ (W/m-K)
ST2212	0.61	0.12	6.9	1.4	0.14
ST2207	1.19	0.22	10.7	2.0	0.20
ST2217	1.74	0.25	12.5	1.8	0.18

## Comparison with Literature TC Values

Previous studies on  $pH_2$  single crystals grown from the gas phase in an enclosed cell near 10 K.

V.G. Manzhelii, B.Ya. Gorodilov, and A.I. Krivchikov, "Heat transfer in solid parahydrogen with heavy impurities (neon, argon)," Low Temp. Phys. v22, p131 (1996).




FIG. 1. Thermal conductivity of crystals of pure  $pH_2$  and  $pH_2$  with Ne impurity (the concentration in ppm are indicated); the solid lines are calculated results.

ST2330a&b  
1100 PPM O<sub>2</sub>/pH<sub>2</sub>  
photolyzed at 193 nm

